

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The Scarlet Ibis (Guara rubra) in Colorado.—My friend Mr. Livesey informs me that whilst he and a party of friends were duck shooting on Grape Creek in Wet Mountain Valley, Custer County, Colorado, in the month of May, 1876, one of the party shot an exceedingly fine specimen of this rare visitor. This bird was skinned and afterwards mounted by a well known firm of taxidermists. So far as I am aware, this specimen, which is now in Mr. Livesey's possession, was never recorded, and hence my reason for making mention of it.—Willoughby P. Lowe, Pueblo, Col.

The Sandhill Crane (Grus mexicana).—A Correction.—In 'The Auk,' Vol. VIII, pp. 308, 309, I mentioned the capture of this bird, but I expressed my doubts in the article as to its positive identification. It proves to be the Little Brown Crane (Grus canadensis) by careful measurements and comparison with a specimen of Grus mexicana from Florida. I can find but one other record of this bird for the Atlantic coast, viz.: Brewster, Auk, Vol. VII, p. 89.—Arthur T. Wayne, Mount Pleasant, S. C.

The Northern Phalarope.—A Correction.—In my article 'Notes on Certain Water Birds in Massachusetts' (Auk, XI, July, 1894), at end of third line from bottom of page 226, read *Northern*, for Red. In last word of article, page 228, read *April*, for May.—Geo. H. MACKAY, *Nantucket*, Mass.

Colinus virginianus cubanensis not a Florida Bird.—Being responsible for the introduction of the Cuban Bob-white into the North American list, I desire to state that its introduction was an error, explainable as follows:—

The only Cuban examples of the genus in the National Museum collection, although not agreeing very well with the original description and colored plate of Ortyx cubanensis Gould, were naturally supposed to be that subspecies. Birds from southwestern Florida proving to be more like these than examples from more northern localities in Florida (including those from Miami, on which Ortyx virginianus var. floridanus Coues was based), in fact practically indistinguishable from them, it seemed necessary to refer them to the Cuban form. Mr. Chapman has since shown, however (Bull. Am. Mus. Nat. Hist., IV, 1893, No. 1, p. 200), that two very distinct forms occur in Cuba, the true Ortyx cubanensis Gould, which is apparently confined to the mountains, and a form which is "typical of the very dark Quail from southern Florida," which latter he is inclined to think may have been introduced into Cuba. Mr. Chapman having kindly allowed me to inspect his Cuban specimens I am able to state that the specimens from southern Florida referred by me to cubanensis are not that bird, but, unless distinguished by a new name, must be referred to floridanus.-ROBERT RIDGWAY, U. S. National Museum, Washington, D. C.